

Name: \_\_\_\_\_ Date: \_\_\_/\_\_\_/\_\_\_ MR: \_\_\_\_\_

## PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

Over the **last 2 weeks**, how often have you been bothered by any of the following problems?  
(Use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3

FOR OFFICE CODING   0   +        +        +         
=Total Score:       

If you checked off **any** problems, how **difficult** have these problems made it for you to do your work, take care of things at home, or get along with other people?

<b>Not difficult at all</b> <input type="checkbox"/>	<b>Somewhat difficult</b> <input type="checkbox"/>	<b>Very difficult</b> <input type="checkbox"/>	<b>Extremely difficult</b> <input type="checkbox"/>
---	---	---	--

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

**CIDI Based Bipolar Disorder Screening Scale**

	YES	NO
<p><i>Euphoria Stem Question:</i></p> <p>1. Some people have periods lasting several days when they feel much more excited and full of energy than usual. Their minds go too fast. They talk a lot. They are very restless or unable to sit still and they sometimes do things that are unusual for them, such as driving too fast or spending too much money. Have you ever had a period like this lasting several days or longer?</p> <p><i>If the answer is YES, skip to Criterion B screening question (3). If the answer is NO, answer the Irritability Stem Question next.</i></p>		
<p><i>Irritability Stem Question:</i></p> <p>2. Have you ever had a period lasting several days or longer when most of the time you were so irritable or grouchy that you either started arguments, shouted at people or hit people?</p> <p><i>If the answer is YES, continue to Criterion B screening question (3). If the answer is NO, don't answer any more questions.</i></p>		
<p><i>Criterion B Screening Question:</i></p> <p>3. People who have episodes like this often have changes in their thinking and behavior at the same time, like being more talkative, needing very little sleep, being very restless, going on buying sprees, and behaving in many ways they would normally think inappropriate. Did you ever have any of these changes during your episodes of being excited and full of energy or very irritable or grouchy?</p> <p><i>If the answer is YES, continue to answer the rest of the questions in this form. If the answer is NO, don't answer any more questions.</i></p>		
<p><i>Criterion B Symptom Questions:</i></p> <p>Think of an episode when you had the largest number of changes like these at the same time. During that episode, which of the following changes did you experience?</p>		
<p>1. Were you so irritable that you either started arguments, shouted at people or hit people?</p> <p><i>This first symptom question should be answered only if the euphoria stem question #1 was answered YES.</i></p>		
2. Did you become so restless or fidgety that you paced up and down or couldn't stand still?		
3. Did you do anything else that wasn't usual for you – like talking about things you would normally keep private, or acting in ways that you would usually find embarrassing?		
4. Did you try to do things that were impossible to do, like taking on large amounts of work?		
5. Did you constantly keep changing your plans or activities?		
6. Did you find it hard to keep your mind on what you were doing?		
7. Did your thoughts seem to jump from one thing to another or race through your head so fast you couldn't keep track of them?		
8. Did you sleep far less than usual and still not get tired or sleepy?		
9. Did you spend so much more money than usual that it caused you to have financial trouble?		

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_/\_\_\_/\_\_\_

## GAD-7

Over the last 2 weeks, how often have you been bothered by the following problems?

Not  
at all

Several  
days

More than  
half the  
days

Nearly  
every day

(Use "✓" to indicate your answer)

1. Feeling nervous, anxious or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	1	2	3

(For office coding: Total Score T \_\_\_\_ = \_\_\_\_ + \_\_\_\_ + \_\_\_\_ )

Severity Range:

\_\_\_ 0-4: Minimum

\_\_\_ 5-9: Mild

\_\_\_ 10-14: Moderate

\_\_\_ 15-21: Severe

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

**COLUMBIA-SUICIDE SEVERITY RATING SCALE**  
**Screener/Recent - Self-Report**

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_/\_\_\_/\_\_\_

Answer Questions 1 and 2	In The Past Month	
	YES	NO
1) <i>Have you wished you were dead or wished you could go to sleep and not wake up?</i>		
2) <i>Have you actually had any thoughts about killing yourself?</i>		
If YES to 2, answer questions 3, 4, 5, and 6. If NO to 2, go directly to question 6		
3) <i>Have you thought about how you might do this?</i>		
4) <i>Have you had any intention of acting on these thoughts of killing yourself, as opposed to you have the thoughts but you definitely would not act on them?</i>		
5) <i>Have you started to work out or worked out the details of how to kill yourself?</i> <i>Do you intend to carry out this plan?</i>		
	In the Past 3 Months	
6) <i>Have you done anything, started to do anything, or prepared to do anything to end your life?</i>  Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.  <i>In your entire lifetime, how many times have you done any of these things?</i>		

Therapist Name: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_/\_\_\_/\_\_\_

**Alcohol Use Disorders Identification Test - AUDIT**

Please select the answer that is most correct for you to each of the following questions.

- 1) How often do you have a drink containing alcohol? (If you answer never, jump to questions 9&10)  
(0) Never (1) Monthly or less (2) 2-4 times a month (3) 2-3 times per week (4) 4 or more times a week
- 2) How many drinks containing alcohol do you have on a typical day when you are drinking?  
(0) 1-2 (1) 3-4 (2) 5-6 (3) 7-8 (4) 10 or more
- 3) How often do you have six or more drinks on one occasion?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 4) How often during the last year have you found that you were not able to stop drinking once you had started?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 5) How often during the last year have you failed to do what was normally expected from you because of drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 6) How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 7) How often during the last year have you had a feeling of guilt or remorse after drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 8) How often during the last year have you been unable to remember what happened the night before because you had been drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 9) Have you or someone else been injured as a result of your drinking?  
(0) No (2) Yes, but not in the last year (3) Yes, during the last year
- 10) Has a relative or friend, or a doctor or other health worker, been concerned about your drinking or suggested you cut down?  
(0) No (2) Yes, but not in the last year (3) Yes, during the last year

Total Score: \_\_\_\_\_

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

MR: \_\_\_\_\_

Patient's Name:		Date:	
<b>Drug Abuse Screening Test—DAST-10</b>			
These Questions Refer to the Past 12 Months			
1	Have you used drugs other than those required for medical reasons?	Yes	No
2	Do you abuse more than one drug at a time?	Yes	No
3	Are you unable to stop using drugs when you want to?	Yes	No
4	Have you ever had blackouts or flashbacks as a result of drug use?	Yes	No
5	Do you ever feel bad or guilty about your drug use?	Yes	No
6	Does your spouse (or parents) ever complain about your involvement with drugs?	Yes	No
7	Have you neglected your family because of your use of drugs?	Yes	No
8	Have you engaged in illegal activities in order to obtain drugs?	Yes	No
9	Have you ever experienced withdrawal symptoms (felt sick) when you stopped taking drugs?	Yes	No
10	Have you had medical problems as a result of your drug use (eg, memory loss, hepatitis, convulsions, bleeding)?	Yes	No

Total Score: \_\_\_\_\_

<b>Guidelines for Interpretation of DAST-10</b>		
Interpretation (Each "Yes" response = 1)		
Score	Degree of Problems Related to Drug Abuse	Suggested Action
0	No problems reported	Encouragement and education
1-2	Low level	Risky behavior – feedback and advice
3-5	Moderate level	Harmful behavior – feedback and counseling; possible referral for specialized assessment
6-8	Substantial level	Intensive assessment and referral

Skinner HA. The Drug Abuse Screening Test. *Addictive Behavior*. 1982;7(4):363-371.

Yudko E, Lozhkina O, Fouts A. A comprehensive review of the psychometric properties of the Drug Abuse Screening Test.

*J Subst Abuse Treatment* 2007;32:189-198

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

### Rosenberg's Self-Esteem Scale (Rosenberg, 1965)

**Instructions:** Below is a list of statements dealing with your general feelings about yourself. Please indicate how strongly you agree or disagree with each statement.

	Strongly Agree	Agree	Disagree	Strongly Disagree
1. I feel that I'm a person of worth, at least on an equal plane with others.				
2. I feel that I have a number of good qualities.				
3. All in all, I am inclined to feel that I am a failure.				
4. I am able to do things as well as most other people.				
5. I feel I do not have much to be proud of.				
6. I take a positive attitude toward myself.				
7. On the whole, I am satisfied with myself.				
8. I wish I could have more respect for myself.				
9. I certainly feel useless at times.				
10. At times I think I am no good at all.				

Total Score: \_\_\_\_\_

**Scoring:** To score the items, assign a value to each of the 10 items as follows:

- For items 1, 2, 4, 6, 7: Strongly Agree=3, Agree=2, Disagree=1, and Strongly Disagree=0.
- For items 3, 5, 8, 9, 10 (which are reversed in valence, and noted with the asterisks\*\* below): Strongly Agree=0, Agree=1, Disagree=2, and Strongly Disagree=3.

The scale ranges from 0-30, with 30 indicating the highest score possible. Higher scores indicate higher self-esteem. Scores below 15 indicate low self-esteem.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_ / \_\_\_\_ / \_\_\_\_

## Satisfaction With Life Scale (SWLS)

*Instructions:* Below are five statements that you may agree or disagree with. Using the 1 - 7 scale below, indicate your agreement with each item by placing the appropriate number on the line preceding that item. Please be open and honest in your responding.

7 - Strongly agree

6 - Agree

5 - Slightly agree

4 - Neither agree nor disagree

3 - Slightly disagree

2 - Disagree

1 - Strongly disagree

\_\_\_\_ In most ways my life is close to my ideal.

\_\_\_\_ The conditions of my life are excellent.

\_\_\_\_ I am satisfied with my life.

\_\_\_\_ So far I have gotten the important things I want in life.

\_\_\_\_ If I could live my life over, I would change almost nothing.

Total Score: \_\_\_\_\_

Severity Range:

31 – 35: Extremely satisfied

26 – 30: Satisfied

21 – 25: Slightly satisfied

20: Neutral

15 – 19: Slightly dissatisfied

10 – 14: Dissatisfied

5 - 9: Extremely dissatisfied

Ed Diener, Robert A. Emmons, Randy J. Larsen and Sharon Griffin as noted in the 1985 article in the *Journal of Personality Assessment*.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_